



Pica, ARFID, and Rumination Disorder Interview – ARFID Questionnaire: Self-report (PARDI-AR-Q-Self)

<i>Client Name</i>	Generic Client	<i>Date administered</i>	19 Aug 2026
<i>Date of birth (age)</i>	1 Jan 1999 (27)	<i>Time taken</i>	1 min 23s
<i>Assessor</i>	Dr Emerson Bartholomew		


Results

	Score	Category
ARFID screen	-	Positive
Body mass index	23.9	Healthy weight

Subscale profile

	Score (0-6)
Concern about aversive consequences	3.3
Lack of interest in eating or food	5
Sensory-based avoidance	4.7


ARFID subscale profile



Severity of impact

	Score (0-6)
Severity of impact	4

Severity of impact





Client Name	Generic Client
--------------------	----------------

Interpretation

The PARDI-AR-Q Self-report was administered on 19 August 2026. The respondent's responses are consistent with possible ARFID, a positive screen, and further assessment by clinical interview may be warranted to confirm or rule out an ARFID diagnosis. Across the three ARFID subscales, lack of interest in eating or food was the most prominent (5/6), with lower scores on sensory-based avoidance (4.7/6) and concern about aversive consequences (3.3/6).

Lack of interest in eating or food (5/6). The respondent's responses indicate low appetite or interest in eating, which may include forgetting to eat, low enjoyment of food, or feeling full early. In particular, the respondent endorsed:

- *forgetting to eat or finding it hard to make time to eat (Response: 6/6)*
- *lack of enjoyment in food or eating (Response: 5/6)*
- *feeling full before finishing, or stopping eating sooner than others (Response: 4/6)*

Sensory-based avoidance (4.7/6). The respondent's responses indicate that sensitivity to the taste, texture, or appearance of food may be contributing to their avoidance or restriction of foods. In particular, the respondent endorsed:

- *sensitivity to taste (Response: 5/6)*
- *sensitivity to texture or consistency (Response: 5/6)*
- *sensitivity to the appearance of food (Response: 4/6)*

Concern about aversive consequences (3.3/6). The respondent's responses indicate avoidance driven by fear of an aversive outcome from eating, such as choking, vomiting, an allergic reaction, or pain. In particular, the respondent endorsed:

- *avoiding eating situations because of such worries (Response: 4/6)*
- *avoiding or restricting food for fear that something bad might happen (Response: 3/6)*
- *physical feelings of panic or anxiety linked to such fears (Response: 3/6)*

The respondent reports that their eating affects their interpersonal and social functioning to a moderate degree (4/6). Body mass index is 23.9 kg/m², in the healthy weight range for adults.

Scoring and Interpretation Information

The PARDI-AR-Q comprises 29 respondent-completed items (items 4–32 of the original instrument) that fall into three blocks:

1. Anthropometric items (items 4-5): height and weight (2 items). Together with age and sex, these produce BMI and growth status.
2. Diagnostic items (items 6 to 21): twelve are scored Yes = 1 / No = 0 and drive the screen, and the other four (items 10, 16, 18, 20) are conditionally displayed follow-ups that are not scored — item 10 is a numeric entry (how much weight has been lost), and items 16, 18, and 20 are free text.
3. Rating items scored 0 to 6 (items 22 to 32, 11 items): includes the severity-of-impact index (items 22 to 23, rated 0 = no difficulty, and 6 = extremely difficult) and the three subscale scores (items 24 to 32; 3 items each).

Recall periods are not fixed and vary by item, spanning current, past-month, and past-3-month



Client Name	Generic Client
--------------------	----------------

timeframes.

PARDI-AR-Q Scoring

The PARDI-AR-Q does not produce a single total score. It is not summed into a total score and instead produces a categorical screening result, three subscale scores, a severity-of-impact score, and body mass index with growth percentiles as clinical context. Higher subscale and severity scores indicate greater symptom intensity.

The screening result (positive/negative possible ARFID) is generated by an algorithm applied to the diagnostic items (6 to 21). A positive screen for possible ARFID requires both:

- A recognised problem with eating or avoidance (item 6 or item 7)
- At least one consequence:
 - significant weight loss or difficulty gaining or maintaining weight (items 8, 9, 11, 12, or 13)
 - nutritional deficiency (item 14)
 - dependence on supplements or enteral feeding (items 15, 17, or 19)
 - psychosocial impairment (item 21).

Because several different combinations of items can produce a positive result, two people who both screen positive may have endorsed quite different items.

A positive screen indicates possible ARFID; it is not a diagnosis, as the PARDI-AR-Q does not evaluate the exclusion criteria for ARFID, it does not assess whether the eating difficulty is better accounted for by another condition, such as anorexia nervosa or bulimia nervosa, or by a concurrent medical condition or other mental disorder.

The subscale scores and the severity-of-impact score are calculated as the average of their respective items, yielding scores that range from 0 to 6:

- Sensory-based avoidance (items 24 to 26): higher scores indicate that avoidance or restriction driven by sensitivity to the taste, texture, or appearance of food is driving avoidance, often presenting as a narrow range of accepted foods and difficulty trying new foods.
- Lack of interest in eating or food (items 27 to 29): higher scores indicate reduced interest in food or eating, including forgetting to eat, low enjoyment of food, or early fullness, often presenting as inadequate intake without active fear or sensory aversion.
- Concern about aversive consequences (items 30 to 32): higher scores indicate avoidance driven by fear of an aversive outcome such as choking, vomiting, allergic reaction, or pain, often presenting as avoidance of specific foods or eating situations linked to those fears.
- Severity of impact (items 22 and 23): higher scores indicate greater disruption to interpersonal relationships and participation in social situations arising from the eating difficulty.

The three subscale and severity scores are reported on their original 0 to 6 metric and interpreted directly against the response scale, which runs from the absence of the feature (0) to its most extreme expression (6).

Response-scale anchors:

- Sensory-based avoidance: 0 = no particular sensitivity; 6 = extremely negative effect on eating



Client Name	Generic Client
--------------------	----------------

- Lack of interest in eating or food: 0 = never; 6 = always
- Concern about aversive consequences: 0 = never; 6 = always
- Severity of impact: 0 = no difficulty; 6 = extreme difficulty

In the interpretive text section, each of the three subscales is given an explanatory paragraph when its score reaches the midpoint of this scale (3 of 6) or higher; lower-scoring subscales are reported as a number only, and severity of impact is always described.

Differentiating ARFID from paediatric feeding disorder (PFD). Because the parent/carer form extends to children as young as four, interpret elevated scores in young children with particular caution. ARFID and PFD overlap and can co-occur, but PFD encompasses feeding difficulties arising from medical, nutritional, oral-motor/swallowing, feeding-skill, or psychosocial factors, whereas ARFID refers to avoidant or restrictive eating that exceeds what would be expected from these factors or warrants separate clinical attention. The PARDI-AR-Q does not comprehensively assess these underlying contributors or determine whether they better account for the presentation. Elevated scores should therefore be interpreted alongside medical, developmental, oral-motor and swallowing, nutritional, feeding-skill, and psychosocial assessment (Goday et al., 2019).

PARDI-AR-Q Growth Context

For respondents aged 14 to 19, body mass index is reported as a percentile and z-score for age and sex; for those aged 20 and over, body mass index is reported against adult categories. Growth percentiles are derived using the United States Centers for Disease Control and Prevention growth reference (Kuczmarski et al., 2002; Flegal & Cole, 2013). These values provide clinical context regarding weight status and growth patterns that may be relevant to ARFID-related consequences such as significant weight loss, failure to achieve expected weight gain, or faltering growth.

PARDI-AR-Q Graphs

On first administration, the three subscales are displayed together on a horizontal bar graph. Additionally, severity of impact is shown on its own horizontal bar graph, divided into five labelled descriptor bands based on the response options. On a repeated administration a line graph with the three subscales is shown, alongside a line graph displaying the severity of impact.

Client Responses



Client Name | Generic Client

Height and Weight

1 Height

171 cm

2 Weight

70 kg

Screening Items

		Yes	No	
3	Do you think you have a problem with eating, involving avoidance or restriction of foods or your eating overall?	1	0	
4	Have other people (for example, doctors, family members, significant others) said that you have a problem with eating, involving avoidance or restriction of foods or your eating overall?	1	0	
5	Have your eating habits led to difficulty maintaining a sufficient weight or, if you are still growing, difficulty gaining enough weight to keep pace with your growth?	1	0	
6	Have your eating habits led to you losing weight (in other words, if you have lost weight, this is because of avoidance or restriction and not because of a medical illness, or other reason)?	1	0	
7	If yes to the previous question, how much weight have you lost in the past 3 months? (please enter numbers)			
	Not Completed			
8	Have others (for example, doctors, family members) been concerned about your weight loss, or been concerned that you are having difficulty gaining enough weight to grow, or having difficulty maintaining your weight due to your eating habits?	1	0	
		Yes	No	I have finished growing
9	Have others (for example, doctors, family members) been concerned that you are not growing taller as you should due to your eating habits?	1	0	0
		Yes	No	
10	Have you ever been told by any health professional that due to your eating habits you were not growing as expected, or that your height was less than it should be?	1	0	
11	Over the past month, has any health professional said that you have a nutritional deficiency due to your eating habits (for example, low iron, low vitamin B12, low vitamin C)?	1	0	
12	Over the past month, has a healthcare professional prescribed special supplements (for example, pills, capsules, powders, or drinks containing vitamins and/or minerals and other micronutrients) specifically to help with your nutrition?	1	0	
13	If yes to the previous question, what have you been prescribed and how much do you take each day?			
	Not Completed			
14	Over the past month, has a healthcare professional prescribed special supplements (for example, high-calorie drinks or 'shots', or dessert-style high-calorie supplements) specifically to help you maintain or gain weight?	1	0	



Client Name Generic Client

Client Responses (cont.)

15	If yes to the previous question, what have you been prescribed and how much do you take each day?		
	Not Completed		
16	Are you currently receiving any tube feeding (receiving food or fluid via a tube in your nose or into your stomach)?	1	0
17	If yes to the previous question, what is the name of the food or fluid product you take via the tube and how much do you take each day?		
	Not Completed		
18	Does your eating cause you difficulties in daily functioning - that is, in how you are able to go about things each day? This might be at school/college/work or when you are at home.	1	0

Severity of Impact

		0	1	2	3	4	5	6
19	Does your eating cause you difficulties in interactions with other people (for example, disagreements or arguments with parents, siblings, significant others, co-workers), or difficulty making or sustaining friendships or other close relationships? Indicate how difficult interactions with other people are for you because of your eating, ranging from 0 (= no difficulty) to 6 (= extremely difficult).	0	1	2	3	4	5	6
20	Does your eating cause you difficulties in social situations, for example does it make it difficult for you to go out with friends, eat at school/college/work, or stay away from home? Indicate how difficult social situations are for you because of your eating, ranging from 0 (= no difficulty) to 6 (= extreme difficulties/try to avoid all social situations).	0	1	2	3	4	5	6

Sensory-based Avoidance

		0	1	2	3	4	5	6
21	Over the past month, have you been particularly sensitive to variation in taste (for example, noticing slight differences in the taste of foods), which has put you off eating any foods or trying any new foods? Indicate how much sensitivity to taste has affected your eating, ranging from 0 (= no negative effect/no particular sensitivity to taste) to 6 (= extremely negative effect, for example, leading to refusal to eat many foods, sticking only to a limited number of preferred foods, or extreme caution when eating).	0	1	2	3	4	5	6
22	Over the past month have you been particularly sensitive to the texture or consistency of food, which has put you off eating any foods or trying any new foods (for example, do you stick to foods of a certain texture only or have you had difficulty eating foods that have different textures mixed together such as pasta with sauce or sandwiches)? Indicate how much sensitivity to texture or consistency has affected your eating, ranging from 0 (= no negative effect/no particular sensitivity) to 6 (= extremely negative effect, for example, leading to refusal to eat many foods, sticking only to a limited number of preferred foods, or extreme caution when eating).	0	1	2	3	4	5	6
23	Over the past month, have you been particularly sensitive to the appearance of food, which has put you off eating any foods or trying any new foods (for example, if food does not look 'right', such as burnt ends of chips/fries, broken biscuits/cookies, or being the 'wrong' colour)? Indicate how much sensitivity to the appearance of food has affected your eating, ranging from 0 (= no negative effect/no particular sensitivity) to 6 (= extremely negative effect, for example, leading to refusal to eat many foods, sticking only to a limited number of preferred foods, or extreme caution when eating).	0	1	2	3	4	5	6

Lack of Interest in Eating or Food

		0	1	2	3	4	5	6
24	Over the past month, how often have you forgotten to eat or found it difficult to make time to eat? Indicate how often you have forgotten to eat or found it difficult to make time to eat, ranging from 0 (= never) to 6 (= always).	0	1	2	3	4	5	6
25	Over the past month, how often have you lacked enjoyment in food or eating (even if only certain foods)? Indicate how often you have lacked enjoyment in food or eating, ranging from 0 (= never) to 6 (= always).	0	1	2	3	4	5	6
26	Over the past month, how often have you felt full before your meal is finished, or stopped eating sooner than others because you had had enough? Indicate how often you have felt full or stopped eating early, ranging from 0 (= never) to 6 (= always).	0	1	2	3	4	5	6



Client Name	Generic Client
--------------------	----------------

Concern about Aversive Consequences

		0	1	2	3	4	5	6
27	Over the past month have you been avoiding or restricting the amount or type of food you eat, because you were afraid that something bad might happen, like being sick, choking, having an allergic reaction, or being in pain? Indicate how often being afraid something bad might happen has affected your eating, ranging from 0 (= never) to 6 (= always).	0	1	2	3	4	5	6
28	Over the past month have you avoided eating situations because you were worried something bad might happen, like being sick, choking, having an allergic reaction, or being in pain while eating (for example, because you might be served something you usually avoid for these reasons, or because you have had a bad experience in the past)? Indicate how often you have avoided eating situations due to such worries, ranging from 0 (= never) to 6 (= always).	0	1	2	3	4	5	6
29	Over the past month have you had any physical feelings of panic or anxiety (examples might include a racing heart, sweaty palms, feeling sick) when you have seen something that has made you think something bad might happen, like being sick, choking, having an allergic reaction, or being in pain while eating. Indicate how often you have had physical feelings of panic or anxiety due to such thoughts, ranging from 0 (= never) to 6 (= always).	0	1	2	3	4	5	6